



Native Fishery Report Cards – 2023:


Report cards for nine important recreational and threatened non-recreational native freshwater fish from 10 priority streams



Energy, Environment and Climate Action



December 2023
Victorian Fisheries Authority
Science Report Series No. 42



Native Fishery Report Cards – 2023:

Report cards for nine important recreational and threatened non-recreational native freshwater fish from 10 priority streams

Brett A. Ingram, Jason Lieschke and Justin Bell

December 2023

**Victorian Fisheries Authority
Science Report Series No. 42**

Published by the Victorian Government, Victorian Fisheries Authority (VFA),
December 2023

© The State of Victoria, Victorian Fisheries Authority Melbourne, 2023

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the *Copyright Act 1968*.

Authorised by the Victorian Government, 1 Spring Street, Melbourne.

Printed by VFA Queenscliff, Victoria.

Preferred way to cite this publication:

Ingram, B.A., Lieschke, J. and Bell, J. (2023). Native Fishery Report Cards – 2023: Report cards nine important recreational and threatened non-recreational native freshwater fish from 10 priority streams. Victorian Fisheries Authority Science Report Series No. 42. 58 pp.

ISSN 2203-3122 (Print)

ISSN 2204-6933 (Online)

ISBN 978-0-6457979-8-5 (Print)

ISBN 978-0-6457979-9-2 (pdf/online/MS Word)

Front cover

Surveying inland rivers with drum nets in the early 1950s (photos courtesy of Mr John Cooper formerly of the Department of Fisheries and Wildlife) compared to electrofishing in the 2020s (photos supplied by the Arthur Rylah Institute for Environmental Research).

Author Contact Details:

Dr. Brett Ingram and Dr Justin Bell

Fisheries Management and Science Branch, Victorian Fisheries Authority
Private Bag 20, Alexandra. Vic. 3714.

Jason Lieschke

Arthur Rylah Institute for Environmental Research
Department of Energy, Environment and Climate Action
123 Brown Street, Heidelberg, Victoria. Vic. 3084.

Copies are available from the website www.vfa.vic.gov.au

For more information contact the VFA Customer Service Centre 136 186

Disclaimer

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

Accessibility

If you would like to receive this publication in an accessible format, such as large print or audio, please telephone 136 186, or email customer.service@vfa.vic.gov.au

Deaf, hearing impaired or speech impaired? Call us via the National Relay Service on 133 677 or visit www.relayservice.com.au

This document is also available in PDF format on the internet at www.vfa.vic.gov.au

Contents

Executive Summary	1
<hr/>	
Introduction	3
Fishery report cards	3
Objectives	3
Materials and Methods	5
Species, species distribution and conservation status	5
Contemporary survey data (2017 to present)	5
Historic survey data (pre-2017)	6
Angler diary records for the Glenelg River estuary perch	6
How to read the <i>Native Fishery Report Cards</i>	6
Overall rating	6
Fish population performance measures (health indicators)	6
Assessment statement	7
Map	7
Monitoring Results	7
Results and discussion	8
Sites surveyed	8
Species present	8
Native fish stockings	8
Fish abundance	9
Fish sizes	10
Native fishery Report Card results	11
Small-bodied native fish	12
Acknowledgements	13
<hr/>	
References	13
<hr/>	
Native Fishery Report Card – 2023: Australian bass	15
<hr/>	
Native Fishery Report Card – 2023: Estuary perch	17
<hr/>	
Native Fishery Report Card – 2023: Freshwater catfish	19
<hr/>	
Native Fishery Report Card – 2023: Golden perch	21
<hr/>	
Native Fishery Report Card – 2023: Macquarie perch	24
<hr/>	
Native Fishery Report Card – 2023: Murray cod	26
<hr/>	
Native Fishery Report Card – 2023: River blackfish	29

Native Fishery Report Card – 2023: Silver perch	31
Native Fishery Report Card – 2023: Trout cod	33
Appendix I: Common and scientific names of fish	35
Appendix II: Performance measures rules and scoring	37
Appendix III: Common species present in 2023 surveys	40
Appendix IV: Abundance of native fish populations	42
Appendix V: Native fish length – weight relationships	44
Appendix VI: Summary of key health indicators for native fish species since 2019	46
Appendix VII: Status of small-bodied native fish in 2023	47
Appendix VIII: A proposed generalised harvest strategy for freshwater native fish of recreational value	53
Background	53
Objective	53
Methodology and decision rules	53
Data source	53
Assessment parameters	54
Decision rules	54
Provisional harvest strategy assessment of Murray cod and golden perch	54
Murray cod	54
Golden perch	54
References	58

List of Tables

Table 1. Priority stream and large-bodied species surveyed for the <i>Native Fishery Report Cards</i>	4
Table 2. Large-bodied species and rivers assessed for each species that are presented in a fishery report card format in this report.	5
Table 3. Species present in rivers surveyed in 2023.....	8
Table 4. Number of native fish stocked into the reaches of waters surveyed over the last 5 seasons.	9
Table 5. Summary of key health indicators for nine native fish species.	11
Table 6. Health ratings for species in each river assessed in this report, and overall rating.	12
Table 7. Detection ratings for small-bodied native fish in the rivers assessed in this report (ratings based on presence/absence in selected streams since 2017).	12

List of Figures

Figure 1. Location of streams surveyed for native fishery report cards in this report.	4
Figure 2. Largest fish caught during electrofishing surveys of native fish populations conducted in 2023.....	10

Executive Summary

Fishery report cards provide a quick and concise means of presenting complex and detailed fish population data in a simplified format for readers to digest. These report cards are being used by the Victorian Fisheries Authority (VFA) to engage and inform anglers about the status and health of important recreational species in the state.

The *Native Fish Report Card Program (NFRCP)*, which has been conducted annually since 2017, uses fish population surveys to produce *Health cards* for nine important recreational and threatened non-recreational large-bodied native freshwater fish from 10 priority streams in which they occur in Victoria. These species are Australian bass, estuary perch, freshwater catfish, golden perch, Macquarie perch, Murray cod, river blackfish, silver perch and trout cod.

Electrofishing and fyke netting in some locations were used to capture fish. These methods are effective sampling tools that can provide a snapshot of the presence and abundance of fish at the time of sampling. But they do not catch all the fish present and often fish are observed but cannot be caught. Therefore, the numbers of fish presented in the *Native Fishery Report Cards* should be considered as an indication of the fish present only.

Results from these annual surveys are combined with results from similar historic fish surveys dating back to the early 1990s. These are then summarised into a *Native Fishery Report Card* to assess long-term trends in catch rates, fish size structure (presence of multiple year classes, mature fish and recent recruits) and to provide a health assessment for each species in selected streams. Information from the *Angler Fishing Diary Program* was also used in assessment of fish in the Glenelg River.

Native Fishery Report Card results

In 2023, an overall rating of Very Good was assigned to one species (Australian bass), Good for five species (golden perch, Macquarie perch, Murray cod, river blackfish and trout cod), Moderate for one species (estuary perch) and Low for two species (freshwater catfish and silver perch).

Since the previous assessment in 2022, the health assessment for three species has changed:

- Australian bass has improved from Good to Very Good
- Macquarie perch has improved from Moderate to Good
- River blackfish has declined from Very Good to Good.

Further information regarding these assessments for 2023 is provided in the report cards below.

Status of small-bodied native fish

There have been considerable declines in the distribution and abundance of many small-bodied native fish, and some are now considered threatened in Victoria. Although small-bodied native fish are also collected as part of fishery surveys to assess species of commercial and/or recreational value, their status is rarely included in fishery assessment reports. Results from current and historic fisheries surveys described in this report were used to assess the status of 24 species of small-bodied native fish, including two-spined blackfish, bony bream, seven galaxias species, Australian grayling, five gudgeon species, two lamprey species, unspotted hardyhead, three pygmy perch species, Murray-Darling rainbow fish, Australian smelt and tupong.

Small-bodied native fish were likely to be either under-represented or absent from surveys described in this report because the survey methods were not designed to target either small-bodied species or the habitats in which they may be more abundant (e.g. smaller river tributaries and backwaters). Some species are threatened, are less often recorded, have either patchy or restricted distributions and/or occur in small numbers. Consequently, status ratings for small-bodied native fish used presence-absence (detection frequency) data in selected streams.

In the streams surveyed in 2023, eight small-bodied native species (including five galaxias) had a detection rating of rare or absent, two occasionally detected, three regularly detected and 11 commonly detected.

Harvest strategy

Outlines the development of a proposed harvest strategy for key recreational species, including Murray cod and golden perch, which incorporates the assessment of four parameters including:

- CPUE of all fish, as an indicator of fish abundance
- CPUE of fish above the slot limit for Murray cod and over the minimum legal size for other species
- Proportion of fish above the slot limit for Murray cod and over the minimum legal size for other species
- Proportion of 1+ year old (10-20 cm), as an indicator of recent recruits.

Provisional assessment of Murray cod and golden perch indicated that neither species met the criteria for further investigation of the status of the fishery.

Species	Rivers	2023 Results						Overall rating
		5-year abundance	10-year abundance	Multiple year classes	Mature fish	Recent recruitment	Maximum size	
Australian bass	Mitchell R., and Thomson R. & Macalister R.	↔	↔	✓	✓	✓*	Some	Very Good
Estuary perch	Glenelg R.	↓	↔	✓	✓	✗	Some	Moderate
Freshwater catfish	Lindsay R. & Murrumbidgee Ck, and Wimmera R.	↔	↔	?	?	?	?	Low
Golden perch	Goulburn R., Gunbower Ck, Lindsay R. & Murrumbidgee Ck, and Wimmera R.	↔	↔	✓	✓	Some*	Some	Good
Macquarie perch	Ovens R. and Yarra R.	↔	↔	✓	✓	Some*	Some	Good
Murray cod	Goulburn R., Gunbower Ck, Ovens R., Lindsay R. & Murrumbidgee Ck	↓	↔	✓	✓	Some*	Some	Good
River blackfish	Gellibrand R. system	↔	?	✓	✓	Some	Some	Good
Silver perch	Gunbower Ck and Wimmera R.	↔	↔	?	?	?	?	Low
Trout cod	Goulburn R. and Ovens R.	↑	↔	✓	✓	Some	Some	Good

See text for explanation of symbols. * May include stocked fish.

Rare or absent	Occasionally detected	Regularly detected	Commonly detected
Galaxias, climbing	Lamprey, shortheaded	Blackfish, two-spined	Bream, bony
Galaxias, flatheaded*	Pygmy perch, southern*	Gudgeon, Cox's*	Galaxias, common
Galaxias, mountain		Lamprey, pouched	Galaxias, ornate
Galaxias, spotted			Grayling, Australian*
Gudgeon, striped			Gudgeon, carp
Pygmy perch, Yarra*			Gudgeon, flathead
Gudgeon, dwarf flathead			Hardyhead, unspotted
Galaxias, obscure			Pygmy perch, Ewen*
			Rainbowfish, Murray-Darling*
			Smelt, Australian
			Tupong

* (red text) Listed as threatened under the the *Flora and Fauna Guarantee Act (1988)* (<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>)

Introduction

Fishery report cards

Fishery report cards provide a quick and concise way of presenting complex and detailed fish population data in a simplified format for readers to digest. These report cards are being used by the Victorian Fisheries Authority (VFA) to engage and inform anglers about the status and health of important recreational species in the state. Fishery report cards have been prepared for popular estuarine fish, trout and freshwater native fish.

Preparation of the *Native Fishery Report Cards* has been a partnership between the Department of Environment, Energy and Climate Action (DEECA), formerly the [Department of Environment, Land, Water and Planning \(DELWP\)](#), the [Victorian Fisheries Authority \(VFA\)](#) and Recreational Fishing License Holders (through Recreational Fishing License Trust [Recreational Fishing Grants Program](#)). The report cards use information from fish population surveys of important recreational and threatened non-recreational large-bodied native freshwater fish from 10 priority streams in Victoria in which they commonly occur (Table 1, Figure 1). The priority rivers were selected in collaboration with catchment management authorities (CMAs), scientists and expert recreational fishers, and are identified as reference rivers for monitoring and assessment in the Victorian *Freshwater Fisheries Management Plan* (Victorian Fisheries Authority 2018).

Native Fishery Report Cards for large-bodied species presented in this report combine information from contemporary electrofishing surveys with historic electrofishing surveys dating back to the early 1990s to assess long-term trends in catch rates, fish size structure (presence of multiple year classes, mature fish and recent recruits) and the current health of nine native fish species (Australian bass, estuary perch, freshwater catfish, golden perch, Macquarie perch, Murray cod, river blackfish, silver perch and trout cod) in 10 priority Victorian streams (Figure 1). The report cards combine and summarise results for one to four rivers for each species (Table 2). Previously *Native Fishery Report Cards* have been published in 2019, 2020 and 2021 (Ingram *et al.* 2019, Ingram and Lieschke 2021, Ingram and Lieschke 2022). DEECA also provides native fish report cards for fish communities in selected rivers (<https://www.ari.vic.gov.au/research/field-techniques-and-monitoring/native-fish-report-card-program>).

Although small-bodied native fish are regularly collected as part of fishery surveys to assess species of commercial and/or recreational value, their status is rarely included in fishery assessment reports. There has been considerable declines in the distribution and abundance of many small-bodied native fish (Lintermans *et al.* 2020) and some are now considered threatened under the Victorian *Flora and Fauna Guarantee Act (1988)* (<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>) (Appendix I). Fishery surveys conducted for preparation of the *Native Fishery Report Cards* may also be used to assess the status of small-bodied native fish populations in selected Victorian streams.

Objectives

To provide fishers and managers a better understanding of the past and current health of Victorian fish species by assessing health of nine large-bodied native fish in 10 streams and produce *Native Fishery Report Cards* for these species. This report also

- Assesses the status of 24 small-bodied native fish species in 10 streams.
- Outlines the development of a harvest strategy for key recreational species, including Murray cod and golden perch.

Table 1. Priority stream and large-bodied species surveyed for the *Native Fishery Report Cards*.

Priority river	Region	Recreational species	Threatened non-recreational species
Gellibrand River system	Corangamite CMA	River blackfish	
Glenelg River	Glenelg Hopkins CMA	Estuary perch	
Lower Goulburn River	Goulburn Broken CMA	Golden perch Murray cod	Silver perch Trout cod
Gunbower Creek	North Central CMA	Golden perch Murray cod	Silver perch Trout cod
Lindsay River & Mullaroo Creek	Mallee CMA	Golden perch Murray cod	Silver perch
Mitchell River	East Gippsland CMA	Australian bass	
Ovens River	North East CMA	Golden perch Murray cod	Macquarie perch Trout cod
Thomson & Macalister rivers	West Gippsland CMA	Australian bass	
Wimmera River	Wimmera CMA	Freshwater catfish Golden perch	Silver perch
Yarra River	Melbourne Water	Macquarie perch Murray cod	




Figure 1. Location of streams surveyed for native fishery report cards in this report.

Table 2. Large-bodied species and rivers assessed for each species that are presented in a fishery report card format in this report.

Species	Mitchell	Thomson & Macalister	Gellibrand	Glenelg	Ovens	Goulburn	Gunbower	Lindsay & Mullaroo	Wimmera	Yarra
Australian bass										
River blackfish*										
Estuary perch										
Macquarie perch*										
Trout cod*										
Murray cod*										
Golden perch										
Silver perch*										
Freshwater catfish*										

* Listed as threatened under the the *Flora and Fauna Guarantee Act (1988)* (<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>)

Materials and Methods

Species, species distribution and conservation status

Scientific names for species described in this report, and their conservation status under the *Flora and Fauna Guarantee Act (1988)* (FFG Act) (<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>), are provided in Appendix I. Species distribution descriptions in Cadwallader and Backhouse (1983), McDowall (1996) and Lintermans (2007) were used to identify rivers to be assessed for each species.

Contemporary survey data (2017 to present)

Contemporary survey data has been collected since 2017 by sampling fish from priority rivers in autumn to avoid the spring periods of peak migration when some native fish undertake long distance movements into or out of the rivers. Surveys conducted at the same time of the year also allows for a more precise comparison between survey years.

Fish were sampled from multiple sites in each priority river, predominantly using electrofishing. Smaller streams, such as the Gellibrand River system, were surveyed with a backpack electrofisher for approximately 90 minutes, while larger streams were surveyed with an electrofishing boat for 60 to 90 minutes. Fyke netting was also used in two rivers where target species that live on the bottom of the river are difficult to catch using electrofishing. These are river blackfish (Gellibrand River system - all years) and freshwater catfish (Wimmera River – prior to 2020).

Electrofishing and fyke netting are effective sampling tools for providing a snapshot of the presence and abundance of fish present at the time of sampling. But they do not catch all the fish present and often fish are observed but cannot be caught. Therefore, the numbers of fish presented in the *Native Fishery Report Cards* should be considered a sample only. There are likely to be many more fish in the waterways than just those recorded. Fish may also move about and populations will fluctuate due to natural variations over time.

Each year, 8-12 sites were surveyed in each stream, and 90 – 3,000 m (typically up to 2,000 m) of stream was surveyed at each site. The length of fish caught were measured and their abundance (number of fish caught per length of stream) was estimated.

Historic survey data (pre-2017)

The contemporary survey *data* were combined with historic electrofishing survey data that were collected within the same river reaches using similar methods to the current surveys to assess long term trends in abundance (fish per 100 m). These historic data were sourced from:

- VFA fish surveys records (Kaiela Fisheries Station – 1982 to 1993 and Snobs Creek – 1989 to 2011). Goulburn River (1996-2011), Gunbower Creek (1993-2007), Lindsay and Mullaroo (1994-1998), Mitchell River (1990-2001), Ovens River (1992-2011), Thomson and Macalister (1989-1999) and Wimmera River (1994-1999).
- The Living Murray Program (Murray-Darling Basin Authority) (<https://www.mdba.gov.au/publications/brochure/living-murray-program>). Gunbower Creek (2008-2017)
- The project, *Integrating fisher-derived and fishery-independent survey data to better understand and manage the Murray Cod fishery in the Murray-Darling Basin* (Fisheries Research and Development Corporation - FRDC Project 2013/022) (Ingram and Raymond 2018). Goulburn and Ovens rivers (2015-2017)
- Monitoring fish stockings in Victoria: 2014 native fish surveys (Ingram *et al.* 2015). Goulburn River and Gunbower Creek (2014).
- Yarra River surveys conducted by DELWP on behalf of Melbourne Water (Tonkin *et al.* 2017). Yarra River (2007-2015)
- Victorian Environmental Flows Monitoring and Assessment Program (VEFMAP) (DELWP 2017a, DELWP 2017b). Glenelg River (2009-2018), Goulburn River (2004-2019), Thomson and Macalister rivers (2005-2016), Wimmera River (2005-2016), Yarra River (2007-2012)
- Sustainable Rivers Audit (SRA) Program (Murray–Darling Basin Authority) and Southern Basins (SB) Program. Data supplied by DELWP (see Lieschke *et al.* 2013a, Lieschke *et al.* 2013b).

Only historic data collected from locations within the same reach of river where contemporary surveys occurred were used in the analyses.

Angler diary records for the Glenelg River estuary perch

In addition, angler catch rate information from the *Angler Fishing Diary Program (AFDP)* was presented for estuary perch in the Glenelg River. The *AFDP*, which commenced in 1997, is the principal means that the VFA uses to monitor recreational fisheries in Victorian estuaries for assessment purposes (Conron *et al.* 2010, Conron and Oliveiro 2016). The *AFDP* provides time-series data on catch rates (as fish per angler hour), catch length-frequency, and catch age-frequency composition for key target species in selected Victorian recreational fisheries, including estuary perch in the Glenelg River which was previously assessed in 2016 (Ingram *et al.* 2016) and 2021 (Ingram *et al.* 2022). Angler catch rate is not intended to be compared directly against electrofishing catch rate but rather provides an additional line-of-evidence for change in abundance over time.

How to read the *Native Fishery Report Cards*

This section provides a brief description of the information presented in the report cards. A more detailed description of the performance measures, how scores were assigned and how the overall rating for each species was determined is provided in Appendix II.

Each report card is broken into several sections. These are:

Overall rating

Overall rating of the health of the species in the selected rivers assessed as part of the report card, which is a synthesis of the fish population performance measures (health indicators). The Overall ratings are:



Fish population performance measures (health indicators)

Information about the abundance (catch rate) of fish and their sizes (length) provide performance measures used to indicate the health of the population. These performance measures are summarised to determine the overall rating for the species. The performance measures are:

Fish abundance

Trend (change) in fish abundance over last five years and last 10 years as indicated by average annual catch rate. Data may be from various sources including electro-fishing surveys and the *Angler Fishing Diary Program (AFDP)*.

Scores: ↑ Increasing ↓ Decreasing. ↔ Stable or variable (up and down)
? Insufficient information to assess.

Fish size (length/age)

Fish size performance measures are based on measurement (fish length) of fish caught in surveys over the last three years. Assessment is conducted only when 60 or more fish are measured in the three years combined, otherwise information to assess fish size is considered insufficient. Fish size categories are:

- Multiple size classes: A wide range of fish sizes present indicates regular successful spawning events and recruitment is occurring in the stream.

Scores: ✓ Wide range of fish size classes present
Some A few fish size classes present
✗ No fish caught or very few fish size classes present
? Insufficient fish measured to assess

- Mature size classes: Mature fish capable of spawning are present in the stream.

Scores: ✓ Good numbers of mature fish present
Some A few mature fish present
✗ No mature fish present
? Insufficient fish measured to assess

- Recent recruitment: Small fish (recruits presumed to be less than one year old) (<10 cm) are present indicates that fish have spawned recently (in last 12 months) in the stream. This may also indicate recent stocking of hatchery-bred fish.

Scores: ✓ Good numbers of small fish present
Some A few small fish present
✗ No small fish present
? Insufficient fish measured to assess

- Maximum size: The presence of fish approaching maximum size indicates sustainable fishing pressure (see Appendix II for maximum size of species).

Scores: ✓ Good numbers of fish approaching maximum size present
Some A few fish approaching maximum size present
✗ No fish approaching maximum size present
? Insufficient fish measured to assess

Assessment statement

Provides a summary health assessment for the species in the selected rivers assessed.

Map

Map showing locations of selected rivers assessed for the species.

Monitoring Results

Monitoring results on the second page of the report card include the following information.

- Average annual catch rate of species in selected rivers based on all available data, and long-term average catch rate. This information was used to assess stock abundance performance measure, i.e. 5-year abundance trend and 10-year abundance trend.
- Size of fish caught in 2023 surveys, including size range of fish measured, percent of fish that were legal size, percent of fish that were mature and percent of fish that were recent recruits.
- Years that the selected rivers have been stocked with hatchery-bred fish.
- Number of hatchery-bred fish stocked into the selected rivers in the last four seasons.

- Length frequency distribution of fish caught during electro-fishing surveys of the selected rivers over last three years and number of fish caught each year in the selected rivers. This information was used to inform fish size performance measures, e.g. presence of recruits, mature fish and fish approaching maximum size.

Results and discussion

Sites surveyed

In 2023, all sites were surveyed as in previous years. However, in the Glenelg River all sites were surveyed except for two upstream sites that were above the range of estuary perch in the river. Instead, two alternate sites in the Glenelg estuary were surveyed. These new sites will be fished in future surveys.

Species present

Forty-three fish species were reported during surveys conducted in 2023, with 8-20 species observed in each river (Table 3). The most common and abundant species was Australian smelt (47% of fish), which was recorded in all 10 rivers and was the most common in five rivers.

The most common and abundant large-bodied fish species was common carp, which represented 77% of large-bodied fish present. The species was observed in nine rivers and was the most common large-bodied fish present in six rivers (Table 3). Further details on the more common species present in each river are provided in Appendix III.

Table 3. Species present in rivers surveyed in 2023.

River	Number of species caught	Most common species (% of total)	Most common large-bodied species (% of total large-bodied species)
Gellibrand River system (including two sites in tributaries, one in Loves Creek and one in Boggy Creek)	8	Common galaxias (36)	Shortfin eel (45)
Glenelg River	20	Gambusia* (38)	Black bream (35)
Lower Goulburn River	12	Australian smelt (43)	Common carp* (66)
Gunbower Creek	11	Unspecked hardyhead (66)	Common carp* (83)
Lindsay River & Mullaroo Creek	11	Common carp* (63)	Common carp* (94)
Mitchell River (including lower Clifton Creek and Wonnangatta River)	17	Australian smelt (77)	Australian bass (27)
Ovens River	11	Australian smelt (37)	Murray cod (33)
Thomson & Macalister rivers	20	Australian smelt (91)	Common carp* (51)
Wimmera River	9	Common carp* (58)	Common carp* (89)
Yarra River	15	Australian smelt (56)	Common carp* (32)
All rivers	43	Australian smelt (47)	Common carp* (77)

* Introduced species

Native fish stockings

The number of native fish stocked over the last 5 seasons into the reaches of waters surveyed is provided in Table 4. Five inland waters and three coastal waters were stocked whereas three waters were not stocked. Species stocked included Australian bass (3 coastal rivers), estuary perch (1 coastal river), golden perch (4 inland rivers), Macquarie perch (1 inland river), Murray cod (4 inland rivers) and silver perch (1 inland river).

Table 4. Number of native fish stocked into the reaches of waters surveyed over the last 5 seasons.

River/creek	Species	2018/19	2019/20	2020/21	2021/22	2022/23	Total
Gellibrand River	Not stocked						
Glenelg River	Not stocked						
Goulburn River	Golden perch	88,150	61,000	60,000	9,000	55,800	273,950
	Murray cod	61,000	40,000	128,000	124,400	61,400	414,800
Gunbower Creek	Golden perch	70,000	70,000	70,000	65,000		275,000
	Murray cod	50,000	50,690	65,000	60,000	60,000	285,690
Lindsay River	Murray cod			27,000	50,000		77,000
Macalister River	Australian bass	7,000	7,000	22,000	23,000	24,000	83,000
Mitchell River	Australian bass	30,000	44,000	60,000	100,000	67,000	301,000
	Estuary perch			50,000			50,000
Mullaroo Creek	Not stocked						
Ovens River	Golden perch	51,000	54,000	50,000	54,000	50,000	259,000
	Macquarie perch	7,510	700	32,000	40,000	10,500	90,710
Thomson River	Australian bass		1,000	1,000	1,000	1,000	4,000
Wimmera River	Golden perch	150,000	80,000	100,000	167,000	169,000	666,000
	Murray cod				10,000	10,000	20,000
	Silver perch	20,000	50,000	50,000	100,000	161,000	381,000
Yarra River	Not stocked						

Fish abundance

In previous reports (Ingram *et al.* 2019, Ingram and Lieschke 2021, Ingram and Lieschke 2022, Ingram and Lieschke 2023), abundance was presented as number of fish caught by electrofishing per 100 m of river (fish/ 100m). However, in this report, abundance is presented as fish caught per electrofishing machine minute (fish/ min) to be consistent with abundance measures used in other fishery reporting programs, such as review of Victorian fish stocks (e.g. Conron *et al.* 2020) and status of Australian fish stocks (<https://www.fish.gov.au/>).

Abundance records (fish/min) for nine native fish species from 10 river systems (Table 2), recorded in electrofishing surveys conducted between 1982 and 2023, were used to assign five levels of fish abundance (see Appendix IV), which were nominally:

Abundance level	Low	Moderate	High	Very high	Exceptional
Electrofishing catch rate (fish/ min)	< 0.05	0.06-0.27	0.27-0.52	0.54 – 0.99	> 1

Historically, abundance has ranged from low (nil to 0.05 fish/min) (48% of records) to exceptional (>1 fish/min) (1% of records) (Appendix IV), the greatest being 1.37 fish/min recorded for Australian bass in the Mitchell River in 2019 followed by 1.35 fish/min for Murray cod in the Ovens River in 2019. Throughout the 1990s and 2000s abundances were generally low to moderate, however, since the mid-2010s high, very high and exceptional abundances have been recorded for some species in some rivers (Appendix IV).

In 2023, abundance was highest for Australian bass in the Mitchell River (1.1 fish/min), which was due to a large number of small fish and many fish across a range of sizes, followed by Murray cod in the Ovens River (0.7 fish/min) (Appendix IV, Figure IV.3). Freshwater catfish, river blackfish and silver perch had the lowest abundances in 2023. The abundance of freshwater catfish continues to remain low as none were caught in the Lindsay-Mullaroo and only four were caught the Wimmera River. The abundance of river blackfish in the Gellibrand system declined to its lowest point

since surveys commenced in 2017. Although the abundance of silver perch remains relatively low, abundances in 2023 were the highest recorded in both Gunbower Creek and the Wimmera River.

Fish sizes

The length – weight relationships for nine native fish are presented in Appendix V. A summary of size ranges of nine native fish species recorded in selected rivers surveyed in 2023 is also provided in Appendix V. The largest fish caught during surveys for each species in 2023, and the river it was caught in, is presented in Figure 2.




Figure 2. Largest fish caught during electrofishing surveys of native fish populations conducted in 2023.

Native fishery Report Card results

Information for nine native fish species and 10 rivers is presented in the report cards and a summary of the key health indicators for the species assessed are provided in Table 5 and Table 6. An overall rating of Very Good was assigned to one species (Australian bass), Good for five species (golden perch, Macquarie perch, Murray cod, river blackfish and trout cod), Moderate for one species (estuary perch) and Low for two species (freshwater catfish and silver perch).

Since the previous assessment in 2022 (Ingram and Lieschke 2023), the health assessment for three species has changed (see Appendix VI):

- Australian bass has improved from Good to Very Good
- Macquarie perch has improved from Moderate to Good
- River blackfish has declined from Very Good to Good.

Further information regarding these assessments is provided in the report cards below.

Table 5. Summary of key health indicators for nine native fish species.

Species	Rivers	2023 Results						Overall rating
		5-year abundance	10-year abundance	Multiple year classes	Mature fish	Recent recruitment	Maximum size	
Australian bass	Mitchell R., and Thomson R. & Macalister R.	↔	↔	✓	✓	✓*	Some	Very Good
Estuary perch	Glenelg R.	↓	↔	✓	✓	✗	Some	Moderate
Freshwater catfish	Lindsay R. & Mullaroo Ck, and Wimmera R.	↔	↔	?	?	?	?	Low
Golden perch	Goulburn R., Gunbower Ck, Lindsay R. & Mullaroo Ck, and Wimmera R.	↔	↔	✓	✓	Some*	Some	Good
Macquarie perch	Ovens R. and Yarra R.	↔	↔	✓	✓	Some*	Some	Good
Murray cod	Goulburn R., Gunbower Ck, Ovens R., Lindsay R. & Mullaroo Ck	↓	↔	✓	✓	Some*	Some	Good
River blackfish	Gellibrand R. system	↔	?	✓	✓	Some	Some	Good
Silver perch	Gunbower Ck and Wimmera R.	↔	↔	?	?	?	?	Low
Trout cod	Goulburn R. and Ovens R.	↑	↔	✓	✓	Some	Some	Good

* May include stocked fish.

Table 6. Health ratings for species in each river assessed in this report, and overall rating.

Species	Mitchell	Thomson & Macalister	Gleneelg	Ovens	Goulburn	Gunbower	Lindsay & Mullaroo	Wimmera	Yarra	Gellibrand	OVERALL
Australian bass	Good	Very good									Very Good
Estuary perch			Moderate								Moderate
Macquarie perch				Good					Very good		Good
Trout cod				Good	Good						Good
Murray cod				Good	Very Good	Good	Good				Good
Golden perch					Very Good	Very good	Good	Good			Good
Silver perch						Low		Low			Low
Freshwater catfish							Low	Moderate			Low
River blackfish										Good	Good

Small-bodied native fish

The status of 24 species of small-bodied native fish were assessed (see Appendix VII). In the streams surveyed, eight species (including five galaxias) had a detection rating of rare or absent, two occasionally detected, three regularly detected and 11 commonly detected.

Table 7. Detection ratings for small-bodied native fish in the rivers assessed in this report (ratings based on presence/absence in selected streams since 2017).

Rare or absent	Occasionally detected	Regularly detected	Commonly detected
Galaxias, climbing	Lamprey, shortheaded	Blackfish, two-spined	Bream, bony
Galaxias, flatheaded*	Pygmy perch, southern*	Gudgeon, Cox's*	Galaxias, common
Galaxias, mountain		Lamprey, pouched	Galaxias, ornate
Galaxias, spotted			Grayling, Australian*
Gudgeon, striped			Gudgeon, carp
Pygmy perch, Yarra*			Gudgeon, flathead
Gudgeon, dwarf flathead			Hardyhead, unspotted
Galaxias, obscure			Pygmy perch, Ewen*
			Rainbowfish, Murray-Darling*
			Smelt, Australian
			Tupong

* Listed as threatened under the the *Flora and Fauna Guarantee Act (1988)*

(<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>)

Acknowledgements

The authors acknowledge the support of the Department of Energy and Climate Action (DEECA), formerly Department of Environment, Land, Water and Planning (DELWP), the Victorian Fisheries Authority (VFA) and the Recreational Fishing License Holders (through Recreational Fishing License Trust Recreational Fishing Grants Program) for funding the *Native Fish Report Card Program* (NFRCP).

The following ARI staff are thanked for assisting with surveys. Lauren Johnson, Adrian Kitchingman, Matthew Jones, Justin O'Connor, John Mahoney, Andrew Pickworth and Frank Amtstaetter.

Dion Iervasi from Austral Ecology sampled the Wimmera.

The authors wish to thank Fiona Warry (DEECA) and Taylor Hunt (VFA) for supporting the Native Fish Report Card Program (NFRCP) and development of these report cards for anglers. Fiona Warry (DEECA), Paul Reich (DEECA), Dr Greg Woodward (DEECA), Dr Taylor Hunt (VFA), John Douglas (VFA), Dr Harry Gorfine (VFA) and Simon Conron (VFA) are also thanked for their advice and comments on the report cards, performance measures and scoring methods therein.

References

- Bayley, P.B. and Austen, D.J. (2002). Capture efficiency of a boat electrofisher. *Transactions of the American Fisheries Society* 131 (3): 435-451.
- Buckmeier, D.L. and Schlechte, J.W. (2009). Capture Efficiency and Size Selectivity of Channel Catfish and Blue Catfish Sampling Gears. *North American Journal of Fisheries Management* 29 (2): 404-416.
- Cadwallader, P.L. and Backhouse, G.N. (1983). *A Guide to the Freshwater Fish of Victoria*. Victorian Government Printing Office, Melbourne. 249 pp.
- Conron, S., Bell, J.D., Ingram, B.A. and Gorfine, H.K. (2020). *Review of key Victorian fish stocks - 2019*. Victorian Fisheries Authority Science Report Series No. 15. Victorian Fisheries Authority, Queenscliff. 173 pp.
- Conron, S., Bridge, N.F., Gixti, D., Ward, M. and Stokie, T.K. (2010). *Angler diary monitoring of recreational target fish stocks in selected Victorian estuaries*. Recreational Fishing Grant Program Final Report (Project numbers: 5/05/06/1, 06/07/3 and 07/08/14). Department of Primary Industries, Victoria, Queenscliff.
- Conron, S.D. and Oliveiro, P. (2016). *State-wide Angler fishing Diary Program 2011-14* Recreational Fishing Grants Program Research Report June 2016. Department of Economic Development, Jobs, Transport and Resources, Queenscliff. 45 pp.
- Cowx, I.G. and Lamarque, P. eds. (1990). *Fishing with Electricity. Applications in Freshwater Fisheries Management*. Fishing News Books. 248 pp.
- Davies, P.E., Harris, J.H., Hillman, T.J. and Walker, K.F. (2010). The sustainable rivers audit: assessing river ecosystem health in the Murray-Darling Basin, Australia. *Marine and Freshwater Research* 61: 764-777.
- DELWP (2017a). *VEFMAP Stage 6 Part A: Program context and rationale*. A report by Arthur Rylah Institute for Environmental Research and Integrated Water and Catchments Division. Department of Environment, Land, Water and Planning, Victoria., Heidelberg.
- DELWP (2017b). *VEFMAP Stage 6 Part B: Program design and monitoring methods* A report by Arthur Rylah Institute for Environmental Research and Integrated Water and Catchments Division. Department of Environment, Land, Water and Planning, Victoria., Heidelberg.
- Ingram, B.A., Hall, K. and Conron, S. (2016). *Recreational fishery assessment 2016 – western estuaries* Recreational Fishing Grants Program Research Report, December 2016. Victorian Government, Department of Economic Development, Jobs, Transport and Resources, Melbourne. 42 pp.
- Ingram, B.A., Hunt, T.L., Lieschke, J. and Douglas, J. (2015). *Monitoring fish stockings in Victoria: 2014 native fish surveys*. Recreational Fishing Grants Program Research Report. Department of Economic Development, Jobs, Transport and Resources, Queenscliff. 50 pp.
- Ingram, B.A. and Lieschke, J. (2021). *Native Fish Report Cards – 2020* Victorian Fisheries Authority Science Report Series No. 16. Victorian Fisheries Authority 33 pp.

Ingram, B.A. and Lieschke, J. (2022). *Native Fishery Report Cards – 2021: Report cards for nine native fish species and status of small-bodied native fish species*. Science Report Series No. 31. Victorian Fisheries Authority. 51 pp.

Ingram, B.A. and Lieschke, J. (2023). *Native Fishery Report Cards – 2022: Report cards nine important recreational and threatened non-recreational native freshwater fish from 10 priority streams*. Victorian Fisheries Authority Science Report Series No. 33. 51 pp.

Ingram, B.A., Lieschke, J. and Warry, F. (2019). Native fish report cards - 2019. Pp. 13-40. In: *2019 Murray Codference: Conference Proceedings (Shepparton, 8th December 2019)*. Victorian Fisheries Authority, Melbourne.

Ingram, B.A., Oliveiro, P. and Conron, S. (2022). *Angler Diary Program: Report cards for key species in coastal recreational fisheries, 2021*. Victorian Fisheries Authority Science Report Series No. 28. 48 pp.

Ingram, B.A. and Raymond, S. (2018). *Implementing better practice Murray cod fishery management*. Recreational Fishing Grants Program Research Report. November 2018. Victorian Fisheries Authority, Melbourne. 50 pp.

Lieschke, J.A., Dodd, L., Stoessel, D., Raadik, T., A., S., Kitchingman, A. and Ramsey, D. (2013a). *The status of fish populations in Victorian rivers 2004–2011 — Part B: Individual basin assessments*. Arthur Rylah Institute for Environmental Research Technical Report Series No. 247. Department of Environment and Primary Industries, Heidelberg, Victoria. 353 pp.

Lieschke, J.A., Dodd, L., Stoessel, D., Raadik, T.A., A., S., Kitchingman, A. and Ramsey, D. (2013b). *The status of fish populations in Victorian rivers 2004–2011 – Part A* Arthur Rylah Institute for Environmental Research Technical Report Series No. 246 Department of Environment and Primary Industries, Heidelberg, Victoria. 147 pp.

Lintermans, M. (2007). *Fishes of the Murray-Darling Basin: An Introductory Guide* Murray-Darling Basin Commission, Canberra. 157 pp.

Lintermans, M., Geyle, H.M., Beatty, S., Brown, C., Ebner, B.C., Freeman, R., Hammer, M.P., Humphreys, W.F., Kennard, M.J. and Kern, P. (2020). Big trouble for little fish: identifying Australian freshwater fishes in imminent risk of extinction. *Pacific Conservation Biology* 26 (4): 365-377.

McDowall, R.M. ed. (1996). *Freshwater Fishes of South-eastern Australia*. Reed Books, Chatswood. 247 pp.

MDBC (2004). *Sustainable Rivers Audit Program*. MDBC Publication 38/04. Murray–Darling Basin Commission, Canberra.

Raadik, T.A. (2014). Fifteen from one: A revision of the *Galaxias olidus* Günther, 1866 complex (Teleostei, Galaxiidae) in south-eastern Australia recognises three previously described taxa and describes 12 new species. *Zootaxa* 3898 (1): 1-198.

Reynolds, J.B. and Koltz, A.L. (1983). Electrofishing. In: *Fisheries Techniques* (L.A., N. and Johnson, D.L. eds.), pp. 147-163. American Fisheries Society, Bethesda, Maryland, USA.

Tonkin, Z., Kearns, J., Fanson, B., Mahoney, J., Ayres, R., Raymond, S., Todd, C. and O'Mahony, J. (2017). *An assessment of Macquarie perch population dynamics in the Yarra River*. Unpublished Client Report for Melbourne Water, June 2017. Arthur Rylah Institute for Environmental Research, Department of Environment, Land, Water and Planning, Heidelberg, Victoria.

Victorian Auditor-General's Office (2018). *Follow Up of Selected 2012–13 and 2013–14 Performance Audits*. Independent assurance report to Parliament. PP no 399, Session 2014–18:23. Victorian Auditor-General's Office (VAGO).

Victorian Fisheries Authority (2018). *Freshwater Fisheries Management Plan 2018*. Victorian Fisheries Authority, Melbourne. 79 pp.

Zalewski, M. and Cowx, I.G. (1990). Factors affecting efficiency of electric fishing. In: *Fishing With Electricity. Applications in Freshwater Fisheries Management* (Cowx, I.G. and Lamarque, P. eds.), pp. 89-111. Fishing News Books, Blackwell Scientific Publications, Ltd., Oxford.



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Australian bass


This report card describes the status of Australian bass in two stream systems, Mitchell River (Mit) (including lower Clifton Creek and Wonnangatta River) and Thomson & Macalister rivers (T-M), in 2023 and trends in population key performance measures that are based on scientific data provided by scientific fishery surveys.

OVERALL RATING - 2023: Very Good


Performance measures (health indicators)	Stream		Status
	Mit	T-M	
Stock abundance			
Trend in abundance over the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys.	↔	↑	↔
Trend in abundance over the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys.	?	↔	↔
Fish size (length/age)			
Wide range of fish size classes present in recent 3 years, indicating regular successful spawning events and recruitment to the population.	✓	✓	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 27 cm in the catch in recent 3 years.	✓	✓	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent 3 years.	✓*	✓*	✓*
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 43 cm) in recent 3 years.	✗	Some	Some
Rating	Good	Very Good	Very Good

↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ✗ = Nil present. * May include stocked fish.

Assessment statement
<p>Australian bass represented 27% (Mitchell) and 23% (Thomson & Macalister) of large-bodied fish caught and observed in 2023. Electrofishing catch rate in the Mitchell was stable over the last 5 years whereas catch rate in the Thomson & Macalister increased over the last 5 years but stable over the last 10 years. Over the last 3 years a wide range of fish sizes, mature fish and recruits were observed in both streams. In 2023, the percentage of legal-size fish was higher in the Thomson & Macalister (49%) than the Mitchell (16%). The presence of many small fish may indicate either recent natural recruitment has occurred or presence of recently stocked fish. Large numbers of small fish were caught in the Mitchell over the last 4 years. Large numbers of small fish were observed in the Thomson & Macalister in some years only, suggesting recent recruitment and/or stockings success has been infrequent. Mature fish were present in both streams. Fish approaching maximum size (a sign of a sustainable fishery) were present in the Thomson & Macalister but not the Mitchell. On this basis the overall rating for Australian bass in 2023 was Very Good, which is an improvement on all previous ratings (Good in all previous years).</p>




Australian bass captured and measured during electro-fishing surveys in 2023	Mitchell	Thomson & Macalister
Size range (cm)	4.8-41	4.1-44
Percent (%) that are legal size (≥ 27 cm)	16	49
Percent (%) that are mature (≥ 27 cm)	16	49
Percent (%) that are recent recruits (< 10 cm)	35	24
Number of fish measured	212	88
Stockings of rivers in recent seasons (1,000s stocked)		
2020/21	60	23
2021/22	100	24
2022/23	67	25



Catch rate

Average catch rate (± s.e.) (black line) and long-term average catch rate (blue line) of Australian bass caught during electro-fishing surveys of two streams. Red triangles = stocking years.



Size distribution

Length frequency distribution of Australian bass caught during electro-fishing surveys of two streams (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Estuary perch

This report card describes the status of the estuary perch in the Glenelg River in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys and recreational fishers (*Angler Fishing Diary Program*).


OVERALL RATING - 2023: **Moderate**

Performance measures (health indicators)	Data source		Status
	Angler	Electro	
Stock abundance			
Trend in abundance the last 5 years as indicated by trend in average catch rate from electro-fishing surveys and from <i>Angler Fishing Diary Program</i>	↓	↔	↓
Trend in abundance the last 10 years as indicated by trend in average catch rate from electro-fishing surveys and from <i>Angler Fishing Diary Program</i>	↓	↔	↔
Fish size (length/age)			
Wide range of fish size classes present in recent 3 years, indicating regular successful spawning events and recruitment to the population.	✓	✓	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 27 cm in the catch in recent 3 years.	✓	✓	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent 3 years.	Some	✗	✗
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 55 cm) in recent 3 years.	Some	✗	Some
Rating	Moderate	Good	Moderate


↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ✗ = Nil present.

Assessment statement

Estuary perch represented 18% of large-bodied fish caught of fish in electrofishing surveys in 2023 and 62% of species angled in 2022. Sites surveyed in 2023 have changed since the previous surveys, which may have influenced the catch rate in the current year. Electrofishing catch rates have been stable of the last 5 and 10 years whereas angling catch rates have declined over the last 5 and 10 years. A wide range of fish sizes, including mature fish, were observed in electrofishing surveys and angler catch over the last 3 years. Fifty percent of fish caught by electrofishing in 2023 were legal size. Some small fish, indicating recent recruitment, were caught by anglers but not by electrofishing. Some fish approaching maximum size (a sign of a sustainable fishery) were caught by anglers but not by electrofishing. On this basis the overall rating for estuary perch in the Glenelg River in 2023 was **Moderate**, which is the same rating for 2022.




Estuary perch captured and measured during an electro-fishing survey of the Glenelg River in 2023 and by angler diarists in 2022	Angler Diary	Electro-fishing
Size range (cm)	14-55	12-50
Percent (%) that are legal size (≥ 27 cm)	90	50
Percent (%) that are mature (≥ 27 cm)	90	50
Percent (%) that are recent recruits (< 10 cm)	2	0
Number of fish measured	164	131
Stockings of river in recent seasons (1,000s stocked): NIL		




Electrofishing catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of estuary perch caught during electro-fishing surveys of the Glenelg River.




Angler catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of estuary perch caught by anglers in the Glenelg River (VFA Angler Diary Program).



Size distribution

Length frequency distribution of estuary perch in the Glenelg River caught by anglers and by electro-fishing (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Freshwater catfish

This report card describes the status of the freshwater catfish in two streams, the Lindsay and Mullaroo rivers (L&M) and Wimmera River (Wim) in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.

OVERALL RATING - 2023: Low


Performance measures (health indicators)	Stream		Status
	L&M	Wim	
Stock abundance			
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	↔
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	?	↔	↔
Fish size (length/age)			
Wide range of fish size classes present in recent 3 years, indicating regular successful spawning events and recruitment to the population.	?	?	?
Mature fish capable of spawning present, as indicated by the presence of fish from 30 cm in the catch in recent 3 years.	?	?	?
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent 3 years.	?	?	?
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 36 cm) in recent 3 years.	?	?	?
Rating	Low	Moderate	Low

↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ***** = Nil present.


Assessment statement

Freshwater catfish represented <0.5% of large-bodied fish caught during electrofishing surveys in both the Lindsay-Mullaroo (L&M) and Wimmera River (Wim) in 2023. Only four fish were caught from the Wimmera in the 2023 surveys. Electrofishing catch rate remains stable over the last 5 years for both streams and was stable but low (below long-term average) over the last 10 years for the Wimmera. There were insufficient fish measured over the last 3 years to assess fish size (length frequency) performance measures. On this basis the overall rating for freshwater catfish in 2023 was **Low** due to the very low abundance of fish in the two streams surveyed. Freshwater catfish were rated as Low in all previous report cards reports.


Freshwater catfish can be taken only from waters within the Wimmera Basin. Taking of freshwater catfish is prohibited in all other Victorian waters.



Freshwater catfish captured and measured during electro-fishing surveys of the Wimmera River in 2023	Lindsay & Mullaroo	Wimmera
Size range (cm)		34-45
Percent (%) that are legal size (≥ 30 cm)	Protected	100
Percent (%) that are mature (≥ 30 cm)	0	100
Percent (%) that are recent recruits (< 10 cm)	0	0
Number of fish measured	0	4
Stockings of river in recent seasons (1,000s stocked): NIL		



Catch rate
Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of freshwater catfish caught during electro-fishing surveys of two streams.



Size distribution

Length frequency distribution of freshwater caught during electro-fishing surveys of two streams (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Golden perch

This report card describes the status of golden perch in four streams, the Goulburn River (Gou), Gunbower Creek (Gun), Lindsay and Mullaroo rivers (L&M) and Wimmera River (Wim), in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.


OVERALL RATING - 2023: **Good**

Performance measures (health indicators)	Stream				Status
	Gou	Gun	L&M	Wim	
Stock abundance					
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↑	↔	↔	↔	↔
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	?	↔	↔
Fish size (length/age)					
Wide range of fish size classes present in recent 3 years, indicating regular successful spawning events and recruitment to the population.	✓	✓	✓	✓	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 30 cm in the catch in recent 3 years.	✓	✓	✓	✓	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent 3 years.	✗	✗*	Some	Some*	Some*
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 50 cm) in recent 3 years.	✓	✓	Some	Some	Some
Rating	Very Good	Very Good	Good	Good	Good


↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ✗ = Nil present. * May include stocked fish.

Assessment statement

Golden perch represented 1-11% of large-bodied fish caught in the four streams surveyed, the Goulburn (Gou), Gunbower (Gun), Lindsay-Mullaroo (L&M) and Wimmera (Wim), in 2023. Electrofishing catch rate over the last 10 years has been stable in three streams (Gou, Gun and Gunbower), and over the last 5 years increased in one stream (Gou) and stable in other streams. Catch rates in recent years are above the long-term average in two streams (Gou and Gun). A wide range of fish sizes, including mature fish, are present in all streams. A high percentage (58-94% of fish caught were legal size. Stocking of hatchery-reared juveniles has occurred in three streams (Gou, Gun and Wim). Some small fish were recorded in only two streams (L&M and Wim). Small fish present in Wim may indicate either recent natural recruitment has occurred or presence of recently stocked fish. Small fish in the L&M, which is not stocked, may be due to recent recruitment. Fish approaching maximum size (a sign of a sustainable fishery) were present in all streams. On this basis the overall rating for golden perch in 2023 was **Good**, which is the same as in 2022 and 2021.




Golden perch captured and measured during electro-fishing surveys in 2023	Goulburn	Gunbower	Lindsay & Mullaroo	Wimmera
Size range (cm)	18-55	25-53	23-51	8-55
Percent (%) that are legal size (≥ 30 cm)	88	94	58	92
Percent (%) that are mature (≥ 30 cm)	88	94	58	92
Percent (%) that are recent recruits (< 10 cm)	0	0	0	2
Number of fish measured	78	32	45	49
Stockings of rivers in recent seasons (1,000s stocked)				
2020/21	60	70	-	100
2021/22	9	65	-	167
2022/23	55.8	0	-	169



Catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of golden perch caught during electro-fishing surveys of four streams. Red triangles = stocking years.



Size distribution

Length frequency distribution of golden perch caught during electro-fishing surveys of four streams (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Macquarie perch

This report card describes the status of Macquarie perch in two streams, the Ovens River (Ove) and Yarra River (Yar), in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.

OVERALL RATING - 2023: **Good**


Performance measures (health indicators)	Stream		Status
	Ove	Yar	
Stock abundance			
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	↔
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	↔
Fish size (length/age)			
Wide range of fish size classes present in recent 3 years, indicating regular successful spawning events and recruitment to the population.	✓	✓	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 30 cm in the catch in recent 3 years.	✓	✓	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent 3 years.	Some*	✗	Some*
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 36 cm) in recent 3 years.	Some	✓	Some
Rating	Good	Very good	Good

⬆ = Increasing, yes and positive. ⬇ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ✗ = Nil present. * May include stocked fish.


Assessment statement

Macquarie perch represented 13-18% of large-bodied fish caught in the two streams surveyed, the Ovens (Ove) and Yarra (Yar) in 2023. Electrofishing catch rates over the last 5 and 10 years have been stable in both streams. The catch rate in the Yarra remains below the long-term average whereas the catch rate in the Ovens is above the long-term average, which may be due to stocking of hatchery-bred fish and translocation of fish from Lake Dartmouth in recent years. A wide range of fish sizes, including mature fish and fish approaching maximum size (a sign of a sustainable fishery), were observed in both streams over the last 3 years. Small fish were caught in the Ovens over the last 3 years, which may be from either recent natural spawnings or recent stocking of hatchery-bred fish, or both. But no small fish were caught in the Yarra in the last 3 years. On this basis the overall rating for Macquarie perch in 2023 was **Good**, which is an improvement on all previous ratings (all were Moderate).

Effective 20 December 2023, the take of Macquarie perch is prohibited in all Victorian waters.




Macquarie perch captured and measured during electro-fishing surveys in 2023	Ovens	Yarra
Size range (cm)	9-37	13-39
Percent (%) that are legal size (≥ 35 cm)	Protected	Protected
Percent (%) that are mature (≥ 30 cm)	22	38
Percent (%) that are recent recruits (< 10 cm)	1	0
Number of fish measured	92	34
Stockings of rivers in recent seasons (1,000s stocked)		
2020/21	32	
2021/22	40	
2022/23	10.5	



Catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of Macquarie perch caught during electro-fishing surveys of two streams. Red triangles = stocking years.



Size distribution

Length frequency distribution of Macquarie perch caught during electro-fishing surveys of two streams (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Murray cod

This report card describes the status of Murray cod in four streams, the Goulburn River (Gou), Gunbower Creek (Gun), Lindsay and Mullaroo rivers (L&M) and Ovens River (Ove), The Goulburn in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.


OVERALL RATING - 2023: Good

Performance measures (health indicators)	Stream				Status
	Gou	Gun	L&M	Ove	
Stock abundance					
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↑	↓	↓	↓	↓
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	?	↔	↔
Fish size (length/age)					
Wide range of fish size classes present in recent 3 years, indicating regular successful spawning events and recruitment to the population.	✓	✓	✓	✓	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 55 cm in the catch in recent 3 years.	✓	✓	✓	✓	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent 3 years.	Some*	Some*	Some*	Some	Some*
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 110 cm) in recent 3 years.	Some	✗	Some	Some	Some
Rating	Very good	Good	Good	Good	Good


↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ✗ = Nil present. * = May include stocked fish.

Assessment statement

Murray cod represented 0.5% to 33% of large-bodied fish caught in the four streams surveyed in 2023. Over the last 5 years electrofishing catch rate has increased in one stream (Gou) but decreased in three streams (Gun, L&M and Ove). However, over the last 10 years catch rate was stable in three streams (Gou, Gun and Ove). There were insufficient data to assess 10-year catch trends in one stream (L&M). A wide range of fish size, including mature fish capable of spawning, were observed in all streams. The percent of fish that were legal size (between 55 & 75 cm) in 2023 ranged from 24% (Gou) to 60% (L&M). Some small fish were present in all streams indicating either recent natural recruitment or recent stocking of hatchery-bred fish (Gou, Gun and L&M). Some fish approaching the maximum size (>110 cm) (a sign of a sustainable fishery) were present in three streams (Gou, L&M and Ove) over the last 3 years. On this basis the overall rating for Murray cod in 2023 was **Good**, which is the same as previous three years.




Murray cod captured and measured during electro-fishing surveys in 2023	Goulburn	Gunbower	Lindsay & Mullaroo	Ovens
Size range (cm)	6-123	33-100	9-114	6-79
Percent (%) that are legal size (between 55 & 75 cm)	24	60	37	34
Percent (%) that are mature (≥ 55 cm)	30	65	47	35
Percent (%) that are recent recruits (< 10 cm)	3	0	5	0
Number of fish measured	96	40	19	175
Stockings of rivers in recent seasons (1,000s stocked)				
2020/21	128	65	27	
2021/22	124.4	60	50	
2022/23	61.4	60	0	



Catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of Murray cod caught during electro-fishing surveys of four streams. Red triangles = stocking years.



Size distribution

Length frequency distribution of Murray cod caught during electro-fishing surveys of four streams (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: River blackfish


This report card describes the status of the river blackfish in the Gellibrand River system (Gel) (including two sites in tributaries, one in Loves Creek and one in Boggy Creek) in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.

OVERALL RATING - 2023:	Good
Performance measures (health indicators)	Gellibrand
Stock abundance	
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	?
Fish size (length/age)	
Wide range of fish size classes present in recent three years, indicating regular successful spawning events and recruitment to the population.	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 12 cm in the catch in recent three years.	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent three years.	Some
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 45 cm) in recent three years.	Some
Rating	Good

↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. * = Nil present.


Assessment statement	
<p>River blackfish represented 19% of large-bodied fish caught in the Gellibrand River system in 2023. River blackfish were more abundant in upstream Gellibrand River sites and particularly the tributary sites in Boggy Creek and Loves Creek. Electrofishing catch rate over the last 5 years was stable. However, this trend is strongly influenced by the two tributary sites which grossly inflate the catch rate (0.05-1.5 fish/min over last 5 years) compared to sites in Gellibrand River proper (0-0.1 fish/min over last 5 years). There is insufficient information to assess trend in catch rate over the last 10 years. A wide range of fish size, including fish that are mature, were present over the last three years. Thirty-three percent of fish caught in 2023 were of legal size (≥ 30 cm). Some small fish were collected over the last three years, indicating recent natural recruitment. Some fish approaching the maximum size (>45 cm) (a sign of a sustainable fishery) were also present. On this basis the overall rating for river blackfish in the Gellibrand River system in 2023 was Good, which is a decline on the 2022 the rating (Very Good).</p>	

River blackfish captured and measured during electro-fishing surveys in 2023	Gellibrand
Size range (cm)	12-41
Percent (%) that are legal size (≥ 30 cm)	33
Percent (%) that are mature (≥ 23 cm)	67
Percent (%) that are recent recruits (< 10 cm)	0
Number of fish measured	15
Stockings of rivers in recent seasons (1,000s stocked):	NIL



Catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of river blackfish caught during electro-fishing surveys of the Gellibrand River system.



Size distribution

Length frequency distribution of river blackfish caught during electro-fishing surveys of the Gellibrand River system (Red numbers = number fish measured. LSL = legal size limit).



Energy,
Environment
and Climate Action



Native Fishery Report Card – 2023: Silver perch

This report card describes the status of the silver perch in two streams, the Gunbower Creek (Gun) and Wimmera River (Wim) in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.

OVERALL RATING - 2023:

Low


Performance measures (health indicators)	Stream		Status
	Gun	Wim	
Stock abundance			
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	↔
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	↔
Fish size (length/age)			
Wide range of fish size classes present in recent three years, indicating regular successful spawning events and recruitment to the population.	?	?	?
Mature fish capable of spawning present, as indicated by the presence of fish from 30 cm in the catch in recent three years.	?	?	?
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent three years.	?	?	?
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 36 cm) in recent three years.	?	?	?
Rating	Low	Low	Low

↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. * = Nil present.


Assessment statement

During 2023 electrofishing surveys seven silver perch were caught in Gunbower Creek and three in the Wimmera River. Electrofishing catch rate has been stable over the last 5 years and 10 years for both streams surveyed. Silver perch abundance in the Wimmera River is low despite regular (annual) stockings of fingerlings. There were insufficient fish measured over the last 3 years to assess fish size (length frequency) performance measures. On this basis the overall rating for silver perch in 2023 was **Low** due to the very low abundance of fish in the two streams surveyed. Silver perch were rated as Low in all previous report cards reports.

The taking of silver perch is prohibited in rivers north of the Great Dividing Range (excluding those in the Wimmera Basin). Silver perch inadvertently caught in these waters must be returned to the water alive and with the least possible injury.




Silver perch captured and measured during electro-fishing surveys in 2023	Gunbower	Wimmera
Size range (cm)	17-46	13-36
Percent (%) that are legal size (≥ 30 cm)	Protected	11
Percent (%) that are mature (≥ 30 cm)	64	11
Percent (%) that are recent recruits (< 10 cm)	0	0
Number of fish measured	14	27
Stockings of rivers in recent seasons (1,000s stocked)		
2020/21		50
2021/22		100
2022/23		161




Catch rate

Average catch rate (\pm s.e.) (black line) and long-term average catch rate (blue line) of silver perch caught during electro-fishing surveys of two streams. Red triangles = stocking years.



Size distribution

Length frequency distribution of silver perch caught during electro-fishing surveys of two streams (Red numbers = number fish measured. LSL = legal size limit).



Energy, Environment and Climate Action



Native Fishery Report Card – 2023: Trout cod

This report card describes the status of trout cod in two streams, the Goulburn River (Gou) and Ovens River (Ove) in 2023 and trends in key population performance measures that are based on scientific data provided by scientific fishery surveys.

OVERALL RATING - 2023: **Good**


Performance measures (health indicators)	Stream		Status
	Gou	Ove	
Stock abundance			
Trend in abundance the last 5 years as indicated by trend in average annual catch rate from electro-fishing surveys	↑	↔	↑
Trend in abundance the last 10 years as indicated by trend in average annual catch rate from electro-fishing surveys	↔	↔	↔
Fish size (length/age)			
Wide range of fish size classes present in recent three years, indicating regular successful spawning events and recruitment to the population.	✓	✓	✓
Mature fish capable of spawning present, as indicated by the presence of fish from 30 cm in the catch in recent three years.	✓	✓	✓
Signs of recent recruitment, as indicated by the presence of fish under 10 cm in recent three years.	Some	Some	Some
Signs of sustainable fishing pressure as indicated by the presence of fish approaching maximum size (≥ 60 cm) in recent three years.	✗	Some	Some
Rating	Good	Good	Good

↑ = Increasing, yes and positive. ↓ = Decreasing, no and negative. ↔ = Stable. ? = Insufficient information.
 ✓ = Good numbers present. **Some** = Some present. ✗ = Nil present.



Assessment statement

In 2023 surveys trout cod represented 8% and 11% of the large-bodied fish caught in the Goulburn River and Ovens River, respectively. Electrofishing catch rate over the last 5 years increased in the Goulburn River and was stable in the Ovens River. Catch rate in both rivers was over the last 10 years has been stable. A wide range of fish sizes and mature fish were observed in both streams. Over the last three years some fish approaching the maximum size (>60 cm) (a sign of a sustainable fishery) were present in the Ovens River only. Some small fish (recruits) were caught in both streams, indicating natural spawning in these streams has occurred in recent years. No stockings of hatchery-bred trout cod fingerlings have occurred in these streams in recent years. On this basis the overall rating for trout cod in 2023 was **Good**, which is the same as previous three years.

Trout cod is a protected species. Taking or possessing trout cod is prohibited, except in Lake Sambell and Lake Kerferd where size and bag limits apply.



Trout cod captured and measured during electro-fishing surveys in 2023	Goulburn	Ovens
Size range (cm)	12 - 51	20 - 55
Percent (%) that are legal size	Protected species	
Percent (%) that are mature (≥ 35 cm)	54	65
Percent (%) that are recent recruits (< 10 cm)	0	0
Number of fish measured	63	58
Stockings of rivers in recent seasons (1,000s stocked): NIL		



Size distribution

Length frequency distribution of trout cod caught during electro-fishing surveys of two streams (Red numbers = number fish measured. LSL = legal size limit).

Appendix I: Common and scientific names of fish

Family	Common name	Scientific name	Comments	FFG status*
Mordaciidae	Shortheaded lamprey	<i>Mordacia mordax</i>		
Geotriidae	Pouched lamprey	<i>Geotria australis</i>		
Anguillidae	Longfin eel	<i>Anguilla reinhardtii</i>		
	Shortfin eel	<i>Anguilla australis australis</i>		
Clupeidae	Bony bream	<i>Nematalosa erebi</i>		
Plotosidae	Freshwater catfish	<i>Tandanus tandanus</i>		Endangered
Prototroctidae	Australian grayling	<i>Prototroctes maraena</i>		Endangered
Retropinnidae	Australian smelt	<i>Retropinna semoni</i>		
Galaxiidae	Climbing galaxias	<i>Galaxias brevipinnis</i>	Broad-finned galaxias	
	Common galaxias	<i>Galaxias maculatus</i>		
	Flatheaded galaxias	<i>Galaxias rostratus</i>		Vulnerable
	Mountain galaxias	<i>Galaxias olidus</i>	Part of <i>olidus</i> species complex	
	Obscure galaxias	<i>Galaxias oliros</i>		
	Ornate galaxias	<i>Galaxias ornatus</i>		
	Spotted galaxias	<i>Galaxias truttaceus</i>		
Artherinidae	Unspecked hardyhead	<i>Craterocephalus stercusmuscarum</i>		
Malanotaeniidae	Murray-Darling rainbowfish	<i>Melanotaenia fluviatilis</i>		Endangered
Percichthyidae	Australian bass	<i>Percalates novemaculeata</i>		
	Estuary perch	<i>Percalates colonorum</i>		
	Golden perch	<i>Macquaria ambigua</i>		
	Macquarie perch	<i>Macquaria australasica</i>		Endangered
	Murray cod	<i>Maccullochella peelii</i>		Endangered
	Trout cod	<i>Maccullochella macquariensis</i>		Endangered
Gadopsidae	River blackfish	<i>Gadopsis marmoratus</i>		Critically Endangered (upper Wannon River form)
	Two-spined blackfish	<i>Gadopsis binspinosus</i>		
Nannopercidae	Ewen pygmy perch	<i>Nannoperca variegata</i>	variegated pygmy perch	Endangered
	Southern pygmy perch	<i>Nannoperca australis</i>		Vulnerable (Murray-Darling lineage)
	Yarra pygmy perch	<i>Nannoperca obscura</i>		Vulnerable
Terapontidae	Silver perch	<i>Bidyanus bidyanus</i>		Endangered
Bovichtidae	Tupong (Congolli)	<i>Pseudaphritis urvillii</i>		
Eleotridae	Carp gudgeon	<i>Hypseleotris</i> Spp	Including western carp gudgeon (<i>H. klunzingeri</i>), Midgley's carp gudgeon (<i>H. sp1</i>) and Lake's carp gudgeon (<i>H. sp2</i>)	

Family	Common name	Scientific name	Comments	FFG status*
	Cox's gudgeon	<i>Gobiomorphus coxii</i>		Endangered
	Dwarf flathead gudgeon	<i>Philypnodon macrostomus</i>		
	Flathead gudgeon	<i>Philypnodon grandiceps</i>		
	Striped gudgeon	<i>Gobiomorphus australis</i>		
Cyprinidae	Common carp	<i>Cyprinus carpio</i>	Introduced species	
Percidae	Redfin perch (English perch)	<i>Perca fluviatilis</i>	Introduced species	
Poeciliidae	Gambusia (mosquitofish)	<i>Gambusia holbrooki</i>	Introduced species	
Salmonidae	Brown trout	<i>Salmo trutta</i>	Introduced species	
	Rainbow trout	<i>Oncorhynchus mykiss</i>	Introduced species	

* Species listed as threatened under the the *Flora and Fauna Guarantee Act (1988)* (<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>)

Appendix II: Performance measures rules and scoring

The following section describes:

- the performance measures used to assess fish species in each river and the rules applied to each performance measure to assign a score.
- How scores for each performance measure are summarized across multiple rivers for each species
- How scores for each performance measure are combined to obtain an overall rating for each species.

Assigning scores to performance measures and summation of scores to determine an overall rating for each species are intended to be objective and based on the available information.

Performance measures

Fish abundance

Trend (change) in fish abundance over time as indicated by average annual catch rate. Data may be from various sources including historic and contemporary electro-fishing surveys (as fish collected per electro-fishing machine minute – fish/min) and the *Angler Fishing Diary Program (AFDP)* (as fish caught per angler per hour). Periods of assessment may include the last five years, last 10 years and long-term (all available records for all years). It is noted that fish abundance can be strongly influenced by a number of variables including timing of surveys around recruitment (and stocking) events and environmental conditions that affect catching efficiency, such as stream flow velocity and turbidity.

Scoring rules:	<ul style="list-style-type: none"> • 5-year trend is assessed only when records are available for least 4 of the past 5 years ($\geq 80\%$ of years). • 10-year trend is assessed only when records are available for at least 8 of the past 10 years ($\geq 80\%$ years). • Long-term trend - assessed only when records are available for $\geq 70\%$ of years from earliest to latest record.
Scores:	<ul style="list-style-type: none"> ↑ Increasing (slope of linear regression line > 0.05) ↓ Decreasing. (slope of linear regression < -0.05) ↔ Stable or variable (up and down) (slope of linear trend between -0.05 and 0.05) ? Insufficient information to assess (as per scoring rules).

Fish size (length/age)

Fish size performance measures are based on measurement (fish length) of fish caught in surveys over the last three years. Assessment is conducted only when 60 or more fish are measured across all three years combined, otherwise information to assess fish size is considered insufficient. Fish size categories are:

Multiple size classes

A wide range of fish sizes present indicates regular successful spawning events and recruitment is occurring in the stream over the last three years. The minimum and maximum recorded length for the target species is identified from all available fishery survey records for that species. The range between these values is then divided into 10 size slots and the proportion of sampled fish in each slot is estimated. Scores are then determined based on the number of slots with fish present.

Scoring rules:	<ul style="list-style-type: none"> • Minimum of 60 fish measured over 3 years. 								
Scores:	<table border="1"> <tr> <td style="text-align: center;">✓</td> <td>Wide range of fish size classes present (fish in at least 5 of 10 size slots).</td> </tr> <tr> <td style="text-align: center;">Some</td> <td>A few fish size classes present (fish present in 2 to 4 of 10 size slots)</td> </tr> <tr> <td style="text-align: center;">✗</td> <td>No fish caught or very few fish size classes present (fish present in 1 of 10 size slots).</td> </tr> <tr> <td style="text-align: center;">?</td> <td>Insufficient fish measured to assess (< 60 fish measured).</td> </tr> </table>	✓	Wide range of fish size classes present (fish in at least 5 of 10 size slots).	Some	A few fish size classes present (fish present in 2 to 4 of 10 size slots)	✗	No fish caught or very few fish size classes present (fish present in 1 of 10 size slots).	?	Insufficient fish measured to assess (< 60 fish measured).
✓	Wide range of fish size classes present (fish in at least 5 of 10 size slots).								
Some	A few fish size classes present (fish present in 2 to 4 of 10 size slots)								
✗	No fish caught or very few fish size classes present (fish present in 1 of 10 size slots).								
?	Insufficient fish measured to assess (< 60 fish measured).								

Mature size classes

Mature fish capable of spawning are present in the stream. Size at maturity is indicative only as this may vary between streams, years and sex of fish for each species. Scores are determined based on the proportion of fish sampled that are mature.

Scoring rules:	<ul style="list-style-type: none"> Minimum of 60 fish measured over 3 years. Size at maturity: Australian bass (≥ 27 cm), estuary perch (≥ 27 cm), freshwater catfish (≥ 30 cm), golden perch (≥ 30 cm), Macquarie perch (≥ 30 cm), Murray cod (≥ 55 cm), silver perch (≥ 30 cm) and trout cod (≥ 30 cm). 	
Scores:	✓	Good numbers of mature fish present (10% or more of fish measured are over mature size).
	Some	A few mature fish present (up to 10% of fish measured are over mature size).
	✗	No mature fish present (No fish measured are over mature size).
	?	Insufficient fish measured to assess (< 60 fish measured).

Recent recruitment

Small fish (recruits presumed to be less than one year old) (<10 cm) present indicates that fish have spawned recently (in last 12 months) in the stream. This may also indicate recent stocking of hatchery-bred fish. Size is indicative only as growth of juveniles may vary between species, streams, time of year sampled and from one year to the next. Note that detection of small fish by electro-fishing can be difficult for some species (notably golden perch). Scores are determined based on the proportion of fish sampled that are small (<10 cm).

Scoring rules:	<ul style="list-style-type: none"> Minimum of 60 fish measured over 3 years. Fish with of length of < 10 cm 	
Scores:	✓	Good numbers of small fish present (10% or more of fish measured are recruits).
	Some	A few small fish present (<10% of fish measured are recruits).
	✗	No small fish present (no recruits measured).
	?	Insufficient fish measured to assess (< 60 fish measured).

Maximum size

The presence of fish approaching maximum size indicates sustainable fishing pressure. Maximum size is indicative only as this may vary between streams and sex of fish for each species. Scores are determined based on the proportion of fish sampled that exceed the maximum.

Scoring rules:	<ul style="list-style-type: none"> Minimum of 60 fish measured 3 years. Maximum size: Australian bass (≥ 43 cm), estuary perch (≥ 55 cm), freshwater catfish (≥ 55 cm), golden perch (≥ 50 cm), Macquarie perch (≥ 36 cm), Murray cod (≥ 110 cm), silver perch (≥ 40 cm) and trout cod (≥ 60 cm). 	
Scores:	✓	Good numbers fish approaching maximum size present (5% or more of fish measured are over maximum size).
	Some	A few fish approaching maximum size present (greater than zero, but <5% of fish measured are over maximum size).
	✗	No fish approaching maximum size present (No fish measured are over maximum size).
	?	Insufficient fish measured to assess (< 60 fish measured).

Summation of preformation measures and overall score determination

Estimating a summarised score for a species for which multiple rivers are assessed uses the median value for each performance measure for each river assessed. For example,

Performance measure		River1	River2	River3	Summary
Trend in abundance over the last 5 years:	Slope	-0.073	0.04	0.558	Median = 0.04
	Score	↓	↔	↑	↔
Mature fish present	Proportion of fish measured	0.04	0.076	0.132	Median = 0.076
	Score	Some	Some	✓	Some

To obtain an overall score, a numerical value is assigned to each performance measure according to the score;

- 3 for ↑ and ✓
- 1 for ↔ and **Some**
- 0 for ↓ and ✗ and ?

The maximum numerical score that can be obtained for either a river or summary of rivers is the number of performance measures assessed multiplied by the maximum value of 3. For example,

6 performance measures multiplied by 3 = Maximum numerical score of 18.

The numerical score for each river assessed for a species is determined by adding the scores for each performance measure together and then dividing the value by the maximum numerical score. For example,


Performance measure	Numerical score	Overall score
Trend in abundance over the last 5 years = ↑	3	10 / 18 (0.556)
Trend in abundance over the last 10 years = ?	0	
Wide range of fish size classes present = ✓	3	
Mature fish present = Some	1	
Small fish under 10 cm present ✓	3	
of fish approaching maximum size present = ✗	0	

The following scale is then applied to assign an overall rating for the species (for either a river or summary of rivers):

Overall score	< 0.1	0.1 to < 0.4	0.4 to < 0.6	0.6 to < 0.9	≥ 0.9
Overall rating	LOW	MODERATE	GOOD	VERY GOOD	EXCELLENT

Appendix III: Common species present in 2023 surveys


The most common species present in each river during 2023 electrofishing surveys, as a percentage of total number of all species present. The most common large-bodied fish species caught and observed in each river during 2023 electrofishing surveys, as a percentage of total number of large-bodied species present (excluding small-bodied species).





Lindsay-Mullaroo

- Australian smelt - 5%
- Bony bream - 18%
- Common carp - 63%
- Goldfish - 5%
- Other - 10%



Lindsay-Mullaroo

- Common carp - 94%
- Golden perch - 1%
- Murray cod - 0%
- Other - 0%
- Redfin - 4%



Mitchell

- Australian bass - 4%
- Australian smelt - 77%
- Common carp - 4%
- Longfin eel - 3%
- Other - 11%



Mitchell

- Australian bass - 27%
- Common carp - 26%
- Longfin eel - 20%
- Other - 12%
- Sand mullet - 14%



Ovens

- Australian smelt - 37%
- Common carp - 18%
- Macquarie perch - 11%
- Murray cod - 21%
- Other - 13%




Ovens

- Common carp - 30%
- Macquarie perch - 18%
- Murray cod - 33%
- Other - 8%
- Trout cod - 11%



Thomson-Macalister

- Australian bass - 1%
- Australian smelt - 91%
- Common carp - 3%
- Other - 3%
- Tupong - 1%




Thomson-Macalister

- Australian bass - 23%
- Common carp - 51%
- Longfin eel - 7%
- Other - 7%
- Redfin - 13%



Wimmera

- Australian smelt - 27%
- Common carp - 58%
- Gambusia - 4%
- Other - 8%
- Redfin - 3%



Wimmera

- Common carp - 89%
- Golden perch - 3%
- Other - 0%
- Redfin - 5%
- Silver perch - 3%



Yarra

- Australian smelt - 56%
- Common carp - 7%
- Common galaxias - 22%
- Other - 9%
- Shortfin eel - 6%



Yarra

- Common carp - 32%
- Macquarie perch - 13%
- Murray cod - 9%
- Other - 6%
- Redfin - 9%
- Shortfin eel - 31%

Appendix IV: Abundance of native fish populations

Abundance records (as fish/min) for nine native fish species recorded in electrofishing surveys conducted between 1982 and 2023 for 10 river systems is presented in Figure IV.1. Frequency distribution of these abundancies is provided in Figure IV.1. Abundance levels, quantile ranges and associated fish abundances are provided in Table IV.1.

Average abundance of native fish species recorded in selected rivers surveyed in 2023 is provided in Figure IV.3.




Figure IV.1. Average abundance records for nine native fish species estimated from historic and contemporary catch electrofishing catch records and associated abundance levels.




Figure IV.2. Frequency distribution native fish abundance (fish/min) for different abundance levels (based on average of all sites within each river each year recorded for electrofishing surveys conducted between 1994 and 2023).

Table IV.1. Abundance levels of native fish and associated quantile ranges, abundance ranges (fish/min) and number of observations (based on average of all sites within each river each year recorded for electrofishing surveys conducted between 1994 and 2023).

Abundance level	Quantile range	Average abundance range (fish/min)	Number of observations	Percent of observations (%)
Low	< 0.05	0 – 0.05	161	48
Moderate	0.05 to < 0.25	0.06 – 0.27	117	35
High	0.25 to < 0.5	0.27 – 0.52	38	11
Very high	0.5 to < 0.99	0.54 – 0.99	15	5
Exceptional	≥ 0.99	1.07 – 1.37	3	1






Figure IV.3. Average abundance of native fish species recorded in selected rivers surveyed in 2023.

Appendix V: Native fish length – weight relationships





Size range (total length) of nine native fish species recorded in selected rivers surveyed in 2023
 (Box = interquartile range with 25th, 50th (median) and 75th percentiles.
 Whisker = largest and smallest value within 1.5 times interquartile range)

Appendix VI: Summary of key health indicators for native fish species since 2019

A summary of key health indicators for nine large-bodied native fish in 10 streams assessed over four years.

Species	Rivers	Results				
		2019	2020	2021	2022	2023
Australian bass	Mitchell R.	Moderate*	Good	Good	Good	Good
	Thomson R. & Macalister R.	Good*	Good	Good	Very good	Very Good
	Overall	Good	Good	Good	Good	Very Good
Estuary perch	Glenelg R.	Good	Good	Good	Moderate	Moderate
Freshwater catfish	Lindsay R. & Mullaroo Ck	Low*	Low	Low	Low	Low
	Wimmera R.	Low*	Low	Low	Low	Low
	Overall	Low	Low	Low	Low	Low
Golden perch	Goulburn R.	Very good*	Good	Good	Good	Very good
	Gunbower Ck	Very good	Moderate	Very good	Very good	Very good
	Lindsay R. & Mullaroo Ck	Moderate*	Moderate	Moderate	Moderate	Good
	Wimmera R.	Good*	Good	Good	Good	Good
	Overall	Good	Moderate	Good	Good	Good
Macquarie perch	Ovens R.	Moderate*	Moderate	Moderate	Moderate	Good
	Yarra R.	Good*	Moderate	Moderate	Moderate	Very Good
	Overall	Moderate	Moderate	Moderate	Moderate	Good
Murray cod	Goulburn R.	Good*	Moderate	Moderate	Good	Very Good
	Gunbower Ck	Very good*	Very good	Very good	Good	Good
	Ovens R	Good*	Good	Good	Good	Good
	Lindsay R. & Mullaroo Ck	Very good*	Very good	Good	Good	Good
	Overall	Very Good	Good	Good	Good	Good
River blackfish	Gellibrand R. system	Low*	Low*	Good	Very Good	Good
Silver perch	Gunbower Ck	Low*	Low	Low	Low	Low
	Wimmera R.	Low*	Low	Low	Low	Low
	Overall	Low	Low	Low	Low	Low
Trout cod	Goulburn R.	Very good*	Good	Good	Good	Good
	Ovens R.	Good*	Good	Good	Good	Good
	Overall	Very Good	Good	Good	Good	Good

* Results not presented in report for the year, instead health indicators are based on retrospective assessment.

References

2019: Ingram *et al.* (2019)
 2020: Ingram and Lieschke (2021)
 2021: Ingram and Lieschke (2022)
 2022: Ingram and Lieschke (2023)
 2023: This report.

Species	Mitchell	Thomson & Macalister	Gellibrand	Glenelg	Ovens	Goulburn	Gunbower	Lindsay & Mullaroo	Wimmera	Yarra
Galaxias, flatheaded*										
Hardyhead, unspecked										
Rainbowfish, Murray-Darling*										
Bream, bony										
Gudgeon, dwarf flathead										
Gudgeon, flatheaded										
Pygmy perch, southern*										
Smelt, Australian										

* Listed as threatened under the the *Flora and Fauna Guarantee Act (1988)*
<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>

Table VII.2. Detection ratings for small-bodied native fish based on presence/absence in selected in the previous 5 years (streams surveyed for each species described in Table VII.1).

Detection rating	Description
RARE OR ABSENT	Present in one stream assessed for the species in < 15% of years, or absent in all streams assessed for the species and absent in the previous 5 years
OCCASIONALLY DETECTED	Present in some streams (< 50%) in some years (< 50%) in those streams assessed for the species in the previous 5 years
REGULARLY DETECTED	Present in most streams (≥ 50%) in some years (< 50%) in those streams assessed for the species in the previous 5 years
COMMONLY DETECTED	Present in most streams (≥ 50%) in most years (≥ 50%) in those streams assessed for the species in the previous 5 years

Detection ratings for small-bodied native fish

Detection ratings for 2023 for these species is presented in Table VII.3. In the streams surveyed, eight species (mostly galaxias) had a detection rating of rare or absent, two occasionally detected, three regularly detected and 11 Commonly detected. Since 2022, the detection rating declined from regularly detected to rare or absent, for one species (obscure galaxias) and increased from rare or absent to regularly detected for one species (pouched lamprey) (Figure VII.1). The presence and absence of 24 small-bodied native fish in surveyed streams since 1980 are presented in Figure VII.2.

Galaxiids

Climbing galaxias, common galaxias and spotted galaxias occur in coastal drainages of Victoria. Common galaxias is widespread and commonly detected in all coastal rivers during surveys and was the most common species recorded in the Gellibrand River system in 2023 (Table 3). Common galaxias recorded in the Wimmera River represent an introduced (translocated) population outside the species natural distribution. Climbing galaxias was rarely detected in surveys and was only recorded in the Gellibrand River system in 2019. Despite having a wide distribution, spotted galaxias was rare and only recorded in the Gellibrand River system in 2018.

Flatheaded galaxias, which is listed as Vulnerable under the *FFG Act*, occurs in rivers north of the Great Dividing Range that are connected to the Murray River but was rare having not been detected in the rivers surveyed since 1984 (Ovens River) and 1986 (Goulburn River).

Mountain galaxias was rare and not recorded in surveys since 2014. Mountain galaxias has been reported from the Ovens and Wimmera rivers only.

Obscure galaxias was detected in both the Ovens and Glenelg rivers but not the Wimmera River while ornate galaxias was regularly detected in the Gellibrand River system and was detected in the Yarra River for the first time since 2014.

Gudgeons

Species of carp gudgeon (incorporating western carp gudgeon, Midgley's carp gudgeon and Lake's carp gudgeon) were commonly detected in all five inland rivers in most years.

Flathead gudgeon and dwarf flathead gudgeon are reported to occur in all rivers surveyed for this report. Flathead gudgeon were commonly detected in most years for all rivers except the Gellibrand River system where it was absent in all years. Dwarf flathead gudgeon are rare and not been reported in the last three years. Previously dwarf flathead gudgeon were occasionally detected in the Thomson, Macalister and Mitchell rivers in some years only.

Cox's gudgeon, which is listed as Endangered under the *FFG Act*, and striped gudgeon occur in coastal rivers of east Victoria. Both species were recorded in the Mitchell River in some years but were absent in surveys of the Thomson & Macalister rivers. Striped gudgeon is rare and has not been recorded since 2018. However, it is noted that distinguishing the two species can be difficult and that striped gudgeon in the Mitchell in 2017 and 2018 were likely Cox's gudgeon.

Other species

Australian grayling, which occurs in coastal drainages of Victoria, was commonly recorded in four streams in most years since 2017.

Australian smelt is one of the more widespread and common fish species in Victorian inland waters. The species was recorded in all rivers and in all years since 2017, and was the most common species recorded in five rivers (Lower Goulburn, Mitchell, Ovens, Thomson & Macalister and Yarra rivers) in 2023 (Table 3).

Bony bream is common in the Lindsay River and Mullaroo Creek system being recorded in every year that the streams were surveyed, and was recorded in most years in Gunbower Creek.

Two-spined blackfish occur in rivers in inland north-east Victoria. The species was present in the Ovens Rivers in the last five out of seven years but was absent in the sites surveyed on the lower Goulburn River, which is not unexpected since two-spined blackfish are not thought to occur downstream of Goulburn Weir where the survey sites are located.

Two species of lamprey occur in Victoria. Pouched lampreys are rarely to regularly detected, and in 2023 was recorded from three out of five rivers being recorded in only one stream in the last seven years. Shortheaded lampreys are occasionally recorded in some years, but in 2023 was recorded in one river (Mitchell).

Unspecked hardyhead were commonly detected in four inland rivers in most years and was the most common species observed in Gunbower Creek in 2023 (Table 3).

Ewen (variegated) pygmy perch occurs in the Glenelg River and, although listed as Endangered under the *FFG Act*, was recorded in surveys every year since 2009.

Southern pygmy perch occurs throughout Victorian coastal and inland waters. However, the species was only recorded from three coastal rivers (Thomson & Macalister, Gellibrand and Glenelg rivers) in recent. The Murray-Darling lineage of southern pygmy perch is listed as Vulnerable under the *FFG Act*. No Southern pygmy perch were detected in surveys of inland rivers.

Yarra pygmy perch occurs in coastal rivers west of Melbourne and is listed as Endangered under the *FFG Act*. The species has not been recorded since 2017 (Glenelg River only).

Murray-Darling rainbowfish occurs in inland rivers that flow into the Murray River (Ovens River, Goulburn River, Gunbower Creek and Lindsay-Mullaroo rivers). Murray-Darling rainbowfish are commonly detected in several inland rivers except for the Ovens River where it was last recorded in 2007. The species is listed as Endangered under the *FFG Act*.

Tupong, which occurs in coastal drainages of Victoria was present in most streams surveyed in most years.

Further information on small-bodied native fish is provided by DEECA in native fish report cards for fish communities in selected rivers (<https://www.ari.vic.gov.au/research/field-techniques-and-monitoring/native-fish-report-card-program>).

Table VII.3. Detection ratings for small-bodied native fish in the rivers assessed in this report (ratings based on presence/absence in the previous 5 years, 2019-2023).

Rare or absent	Occasionally detected	Regularly detected	Commonly detected
Galaxias, climbing	Lamprey, shortheaded	Blackfish, two-spined	Bream, bony
Galaxias, flatheaded*	Pygmy perch, southern*	Gudgeon, Cox's*	Galaxias, common
Galaxias, mountain		Lamprey, pouched	Galaxias, ornate
Galaxias, spotted			Grayling, Australian*
Gudgeon, striped			Gudgeon, carp
Pygmy perch, Yarra*			Gudgeon, flathead
Gudgeon, dwarf flathead			Hardyhead, unspocked
Galaxias, obscure			Pygmy perch, Ewen*
			Rainbowfish, Murray-Darling*
			Smelt, Australian
			Tupong

* Listed as threatened under the the *Flora and Fauna Guarantee Act (1988)* (<https://www.environment.vic.gov.au/conserving-threatened-species/threatened-list>)




Figure VII.1. Change in Detection ratings for small-bodied native fish (ratings based on presence/absence in selected streams for previous 5 years)




Figure VII.2. Presence and absence of small-bodied native fish in surveyed streams since 1980 (Each tile represents a year that each stream was surveyed. N = Number of years present / number of years surveyed for all streams).




Figure VII.2. Continued. Presence and absence of small-bodied native fish in surveyed streams since 1980 (Each tile represents a year that each stream was surveyed. N = Number of years present / number of years surveyed for all streams).

Appendix VIII: A proposed generalised harvest strategy for freshwater native fish of recreational value

Background

Harvest strategies are traditionally a commercial fisheries management measure that provide a framework specifying the pre-determined management actions in a fishery to achieve the agreed management objectives.

To be meaningful, harvest strategies require long-term, consistent, reliable, and robust monitoring data, which is widely acknowledged as being difficult to obtain in recreational fisheries that are 'data-poor' and exist in numerous discrete stocks.

Despite this challenge, there has been a push to explore the utility of harvest strategies in recreational fisheries.

In 2018, the Victorian Fisheries Authority committed to 'develop a harvest strategy by 2023 as a separate document and include predetermined triggers for high-risk fisheries after running the Native Fish Report Card Program for five years.' (Victorian Auditor-General's Office 2018).

The Victorian Freshwater Fisheries Management Plan also list a priority to: 'Developing a harvest strategy for selected freshwater fisheries, using information from fish population health monitoring and recreational fisher feedback' (Victorian Fisheries Authority 2018).

It's important that the harvest strategy is considered in context alongside the Native Fish Report Card Program (NFRCP) data and interpretations (including limitations), thus this appendix meets the above commitment and priority for a harvest strategy for select and high-risk freshwater fisheries.

Objective

This report aims to develop a generalised harvest strategy for freshwater native fish of recreational value in Victoria. Since the strategy utilises electro-fishing surveys conducted as part of native fish report cards (Ingram *et al.* 2019, Ingram and Lieschke 2021, Ingram and Lieschke 2022, Ingram and Lieschke 2023), outputs are limited to a small number of rivers (stocked and non-stocked) across the state. Native fish fisheries in lakes are excluded from the strategy due to lack of monitoring data. While the strategy focuses on Murray cod river fisheries, its design will allow for assessment of other native fish fisheries in rivers that are also surveyed for native fish report cards, such as golden perch. The strategy aims to provide trigger point(s) for further investigation rather than defining what action to be taken.

Methodology and decision rules

Data source

The strategy uses electrofishing survey data collected as part of the NFRCP since 2017 (details are provided in the Materials and Methods Section of this report). The catch per unit effort (CPUE) is the nominal catch rate presented as fish per electrofishing minute (fish/min). However, future refinement of the strategy may consider a standardised CPUE (i.e. adjusted for sized-based capture efficiency for each species). There are no commercial fishing records and recreational fishing timeseries data is limited and patchy.

Electro-fishing is a common and well-established sampling method in freshwaters and is reported to be capable of detecting more species and wider range of fish sizes within species than traditional sampling techniques, such as fyke and mesh netting and seining (Reynolds and Koltz 1983, Cowx and Lamarque 1990). However, a weakness of electrofishing is that capture efficiency can be low and variable (e.g. Zalewski and Cowx 1990, Bayley and Austen 2002, Buckmeier and Schlechte 2009). Electrofishing is less effective in openwaters, and large mobile pelagic species that can avoid approaching vessels are especially difficult to catch. Other factors affecting capture efficiency relate to visibility of fish to operators including high currents, high turbidity and increased wave action. Also, electro-fishers cannot operate on rainy days for operator safety reasons. These factors, which can vary between sites surveyed and from day-to-day, are not accounted for in the strategy. Future development and refinement of the harvest strategy may consider excluding sites affected by these factors. For example, the Gunbower Creek Moss Road Site and Three-Quarter Hole Site, which experience very high flow variability associated with discharge from the weir upstream of the sites and the Lindsay River sites which also have high flow variability.

Electrofishing survey sampling methods have changed over time, which limits comparison with historical data. For example, the Sustainable Rivers Audit (SRA) sampling method samples sites chosen using a stratified-random procedure (MDBC 2004, Davies *et al.* 2010) whereas targeted survey sampling concentrates samples in sites where fish are likely to occur. The strategy only uses electrofishing survey data collected as part of the NFRCP, which commenced in 2017.

Nine of the 12 rivers surveyed for the NFRCP are regularly stocked with hatchery-bred fish, including Australian bass, estuary perch, golden perch, Macquarie perch, Murray cod, and silver perch (see table 4 in this report). The presence of stocked fish in these rivers may be masking natural recruitment. Currently the NFRCP does not attempt to distinguish between stocked and naturally recruited fish.

Assessment parameters

Four assessment parameters are used to assess the status of the fishery in each river surveyed:

- *CPUE of all fish*, as an indicator of fish abundance
- *CPUE of fish above the slot limit for Murray cod and over the minimum legal size for other species*
- *Proportion of fish above the slot limit for Murray cod and over the minimum legal size for other species*
- *Proportion of 1+ year old (10-20 cm)*, as an indicator of recent recruits.

For each assessment parameter, three reference lines are proposed:

- *Green line* - Average CPUE for each river surveyed over a 7-year period, from 2017 (the first year of the NFRCP) to 2022.
- *Amber line* – Nominally 40% of green line.
- *Red line* – Pre-2000 average CPUE.

Localised extreme events at the time of sampling, such as flooding, can have a profound effect on CPUE. Consequently, reference lines span multiples years to reduce the impact of these events.

Presence of fish above the slot limit for Murray cod and over the minimum legal size for other species are used as an indicator of mature fish capable of spawning

Presence 1+ year old (10-20 cm) fish is used as an indicator of recent recruitment. Fish of a smaller size are excluded due to the uncertainty of their detection by electrofishing.

Decision rules

The following decision rules are proposed to trigger further investigation into the status of the state recreational fishery for the species being assessed.

1. Assessment is based on the latest two years of the NFRCP electrofishing survey data to account for variations between years
2. Each assessment parameter is applied to each river surveyed
3. The decision to undertake further investigation is triggered by 50% or more of the rivers surveyed being below the 40% of 2017-2022 reference line (amber line) for at least two assessment parameters for the latest two consecutive years.

Provisional harvest strategy assessment of Murray cod and golden perch

Murray cod

Key points

- Four rivers were assessed, the Goulburn River, Gunbower Creek, Lindsay River and Murrumbidgee River, and Ovens River (Table VIII.1 and Figure VIII.1).
- No river was below the trigger level. That is, no river had at least two assessment parameters below 40% of the 2017-2022 reference line for the latest two consecutive years of survey results.
- Assessment did not meet the criteria for further investigation since 50% or more of the rivers assessed were above the trigger level.

Golden perch

Key points

- Four rivers were assessed, the Goulburn River, Gunbower Creek, Lindsay River and Murrumbidgee River, and Wimmera River (Table VIII.2 and Figure VIII.2).
- No river was below the trigger level. That is, no river had at least two assessment parameters below 40% of the 2017-2022 reference line for the latest two consecutive years of survey results.
- Assessment did not meet the criteria for further investigation since 50% or more of the rivers assessed were above the trigger level.

Table VIII.1. Murray cod harvest strategy assessment.

Assessment parameter	Assessment parameter details	Goulburn	Gunbower	Lindsay-Mullaroo	Ovens
CPUE	Number of years parameter under 40% of 2017-2022 CPUE	0	0	1	0
CPUE of fish over legal size	Number of years parameter under 40% of 2017-2022 CPUE	0	0	0	0
Proportion of fish over legal size	Number of years parameter under 40% of 2017-2022 proportion	0	0	0	0
Proportion fish between 10-20 cm	Number of years parameter under 40% of 2017-2022 proportion	1	1	1	1
Trigger level	At least two assessment parameters below 40% of 2017-2022 reference line for the latest two consecutive years	No	No	No	No
	Further investigation of fishery recommended when 50% or more of waters at trigger level	NO			

Table VIII.2. Golden perch harvest strategy assessment.

Assessment parameter	Assessment parameter details	Goulburn	Gunbower	Lindsay-Mullaroo	Wimmera
CPUE	Number of years parameter under 40% of 2017-2022 CPUE	0	0	1	1
CPUE of fish over legal size	Number of years parameter under 40% of 2017-2022 CPUE	0	0	1	0
Proportion of fish over legal size	Number of years parameter under 40% of 2017-2022 proportion	0	0	0	0
Proportion fish between 10-20 cm	Number of years parameter under 40% of 2017-2022 proportion	1	2	2	0
Trigger level	At least two assessment parameters below 40% of 2017-2022 reference line for the latest two consecutive years	No	No	No	No
	Further investigation of fishery recommended when 50% or more of waters at trigger level	NO			




Figure VIII.1. Murray cod harvest strategy assessment parameters (green reference line = Average CPUE for period 2017-2022. Amber line = 40% of average CPUE for period 2017-2022. Red line = Pre-2000 average).




Figure VIII.2. Golden perch harvest strategy assessment parameters (green reference line = Average CPUE for period 2017-2022. Amber line = 40% of average CPUE for period 2017-2022. Red line = Pre-2000 average).

References

- Bayley, P.B. and Austen, D.J. (2002). Capture efficiency of a boat electrofisher. *Transactions of the American Fisheries Society* 131 (3): 435-451.
- Buckmeier, D.L. and Schlechte, J.W. (2009). Capture Efficiency and Size Selectivity of Channel Catfish and Blue Catfish Sampling Gears. *North American Journal of Fisheries Management* 29 (2): 404-416.
- Cowx, I.G. and Lamarque, P. eds. (1990). *Fishing with Electricity. Applications in Freshwater Fisheries Management*. Fishing News Books. 248 pp.
- Davies, P.E., Harris, J.H., Hillman, T.J. and Walker, K.F. (2010). The sustainable rivers audit: assessing river ecosystem health in the Murray-Darling Basin, Australia. *Marine and Freshwater Research* 61: 764-777.
- Ingram, B.A. and Lieschke, J. (2021). *Native Fish Report Cards – 2020* Victorian Fisheries Authority Science Report Series No. 16. Victorian Fisheries Authority 33 pp.
- Ingram, B.A. and Lieschke, J. (2022). *Native Fishery Report Cards – 2021: Report cards for nine native fish species and status of small-bodied native fish species*. Science Report Series No. 31. Victorian Fisheries Authority. 51 pp.
- Ingram, B.A. and Lieschke, J. (2023). *Native Fishery Report Cards – 2022: Report cards nine important recreational and threatened non-recreational native freshwater fish from 10 priority streams*. Victorian Fisheries Authority Science Report Series No. 33. 51 pp.
- Ingram, B.A., Lieschke, J. and Warry, F. (2019). Native fish report cards - 2019. Pp. 13-40. In: *2019 Murray Codference: Conference Proceedings (Shepparton, 8th December 2019)*. Victorian Fisheries Authority, Melbourne.
- MDBC (2004). *Sustainable Rivers Audit Program*. MDBC Publication 38/04. Murray–Darling Basin Commission, Canberra.
- Reynolds, J.B. and Koltz, A.L. (1983). Electrofishing. In: *Fisheries Techniques* (L.A., N. and Johnson, D.L. eds.), pp. 147-163. American Fisheries Society, Bethesda, Maryland, USA.
- Victorian Auditor-General's Office (2018). *Follow Up of Selected 2012–13 and 2013–14 Performance Audits*. Independent assurance report to Parliament. PP no 399, Session 2014–18:23. Victorian Auditor-General's Office (VAGO).
- Victorian Fisheries Authority (2018). *Freshwater Fisheries Management Plan 2018*. Victorian Fisheries Authority, Melbourne. 79 pp.
- Zalewski, M. and Cowx, I.G. (1990). Factors affecting efficiency of electric fishing. In: *Fishing With Electricity. Applications in Freshwater Fisheries Management* (Cowx, I.G. and Lamarque, P. eds.), pp. 89-111. Fishing News Books, Blackwell Scientific Publications, Ltd., Oxford.

Customer Service Centre **136 186**

vfa.vic.gov.au